



Figure 1.

Vaccine concept



Impact of neuraminidase on MLR-chemokine/cytokine production



Optimal formula for efficient in vivo propagation of mature DC carrying phagocytosed antigens




Figure 4.




Figure 5.




Figure 6.

Allgeneic MLR: Chemokine/cytokine production MIP-1 alpha **RANTES** IL-1 beta IL-6 TNF-alpha IFN-gamma 500 pg/mL monocyteS (control)

Figure 7.






Figure 8

 $\mathbb{S}/1$.




Figure 9

10011





Figure 10





11771



Impact of anti-CD43 on monocyte chemokine produktion

Figure 11

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
□ OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.